

Chinese Cabbage/Napa 中国大白菜 (绍菜)

(*Brassica pekinensis*)

CC1373

CR Mini Top F1 *New 2015!*

韩国抗根肿病娃娃菜—春秋娃娃王(杂交种)



娃娃菜 Wa Wa Tsai @ 35~40 days



小绍菜 Mini Napa @ 45~50 days

CHARACTERISTICS 特性

- Mini-type
- 35~40 days for Wa Wa Tsai after transplanting
- 45~50 days for Mini Napa after transplanting
- Excellent shape, compact and uniform
- Dark green outer leaves. Yellow internal color, good texture
- Slow bolting, CR and DM resistant
- For spring, summer and fall
- Planting @ 10" for Wa Wa Tsai; 12" for Mini Napa
- 小型品种, 娃娃菜和小绍菜专用
- 娃娃菜 35~40 天 (移栽后)
- 小绍菜 45~50 天 (移栽后)。3~4 磅
- 球形美观, 紧凑, 大小一直
- 叶色深绿, 内叶淡黄色, 风味品质佳
- 耐抽薹, 抗根肿病和霜霉病
- 适合春、夏、秋栽培
- 栽培密度: 娃娃菜 10"; 小绍菜 12"

• The description is based upon our trials in Lynden Ontario Canada. Customers in other areas are suggested to have trials in different seasons before commercial growing.

• 品种描述基于我们在加拿大安省 Lynden 的试种结果, 其他地区的用户在规模种植之前, 请根据当地情况在不同季节先小规模试种。